

Exploring the Potential of Spirulina (*Arthrospira platensis*) Aqueous Extract in Preventing Glycation of Hemoglobin and pBR322 Plasmid

Pharmacognosy Magazine
19(3) 581–591, 2023
© The Author(s) 2023
Article reuse guidelines:
in.sagepub.com/journals-permissions-india
DOI: 10.1177/09731296231170959
journals.sagepub.com/home/phm



Additiya Paramanya¹, Mohammad Abul Farah², Khalid Mashay Al-Anazi², Hari Prasad Devkota^{3,4} and Ahmad Ali¹ 

Abstract

Background: Spirulina (*Arthrospira platensis*), a cyanobacterium, is being consumed worldwide owing to its high nutritional value and therapeutic potential.

Objectives: This study aims to determine the phytochemical content, antioxidant capacity, and antiglycation property of *A. platensis* PCC 7345 aqueous extract against glucose, fructose, and ribose-mediated glycation of hemoglobin.

Materials and Methods: The antioxidant property of the extract was determined by 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), and a nitric oxide scavenging assay. Protein glycation and its subsequent prevention were interpreted by observing a reduction in browning, fructosamine, glycated hemoglobin (HbA_{1c}) level, protein carbonyl content, advanced glycation end products (AGEs) formation, and protein aggregation using spectrophotometry and spectrofluorometry. The prevention of strand nicking/breaks in the pBR322 plasmid was analyzed using electrophoresis, and the change in integrated density was calculated using ImageJ software.

Results: The extract showed very good antioxidant potential, which is evident and could be estimated from the inhibition concentration (IC₅₀) values that were 45.97, 31.09, and 52.29 µg/mL in the DPPH, ABTS, and nitric oxide scavenging assays, respectively. *A. platensis* aqueous extract was found to be the most effective in preventing glucose-mediated glycation and protein aggregation, wherein 100 µg/mL of the extract could effectively cause ~50% inhibition in the formation of early and AGEs and prevented the formation of protein aggregates. The highest antiglycating potential of the extract was observed in glucose-mediated glycation, followed by fructose and ribose. The formation of four fluorescent AGEs, such as argpyrimidine, vesper lysine A, pentosidine, and crossline, was reduced. The extract could prevent 97% of DNA nicking in pBR322.

Conclusion: The aqueous extract of *A. platensis* has a potent ability to prevent glycation and its secondary complications.

Keywords

Advanced glycation end products (AGEs), *Arthrospira platensis*, glycated hemoglobin, integrated density, protein aggregation

Received 4 November 2022; accepted 1 March 2023

Introduction

Diabetes and its secondary complications are major metabolic disorders affecting millions worldwide. Since the last few decades, there has been a rapid increase in the morbidity and mortality rate of diabetes. Thus, the American Diabetes Association (ADA) recommends screening for prediabetes and type 2 diabetes from the age of 35, even in the absence of any symptoms (American Diabetes Association, 2022). The screening includes determining glycated hemoglobin (HbA_{1c}) levels in the blood. HbA_{1c} is considered to be standard in the estimation of developing long-term complications. Since the half-life of hemoglobin (HB) ranges from 90 to 120 days, the HbA_{1c} level is used as an indicator of glucose control for 2–3 months (American

Diabetes Association, 2022; Singh et al., 2014). On the contrary, fructosamine levels help estimate glucose levels in 7–21 days. This helps in monitoring rapid insulin titrations

¹Department of Life Sciences, University of Mumbai, Mumbai, Maharashtra, India

²Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia

³Graduate School of Pharmaceutical Sciences, Kumamoto University, Kumamoto City, Kumamoto, Japan

⁴Pharmacy Program, Gandaki University, Pokhara, Pokhara, Gandaki Province, Nepal

Corresponding author:

Ahmad Ali, Department of Life Sciences, University of Mumbai, Mumbai 400098, Maharashtra, India.

E-mail: ahmadali@mu.ac.in



and is helpful in the short-term monitoring of diabetes control.

Although not entirely determined, a complex cascade of reactions is activated in the presence of surplus sugar in the body. One of these reactions is glycation, or the non-enzymatic binding of sugar with biomolecules like proteins, lipids, and nucleic acids. Advanced glycation end products (AGEs) formed by protein glycation lead to structural and functional changes, thereby triggering secondary complications of diabetes (Singh et al., 2014).

According to Bunn et al. (1975), the *N*-terminal valine of the β -subunit of HB is the most prone to glycation. It reacts with the carbonyl group of reducing sugars to generate a Schiff base, which further rearranges to form Amadori products. Both products in the initial stages of glycation are formed by a reversible reaction (Thornalley & Rabbani, 2014); hence, these initial steps become the target for antiglycation activity. However, both the products are highly unstable and consequently and parallelly lead to the formation of irreversible AGEs. Though oxidation is not the only process that alleviates AGE formation, a combination of glycation and oxidation, i.e., glycooxidation, further triggers the formation of many AGEs (Ahmad et al., 2014; Thornalley & Rabbani, 2014). These AGEs could be classified into two types based on their ability to emit fluorescence, namely, fluorescent and nonfluorescent AGEs. Fluorescent AGEs get easily detected by simple fluorometric techniques (Thornalley & Rabbani, 2014) and are often used as a marker for glycooxidative damage (Ahmad et al., 2014).

The intermediate carbonyl species originating due to glycation and oxidative stress are highly reactive. They are further known to cause structural and functional changes in other biomacromolecules like nucleic acids and lipids. Glyoxal and methylglyoxal (MG) are the reactive carbonyl species that have been identified as being exclusively responsible for glycation in DNA (Ahmad et al., 2014). These have been used to trigger glycation processes for *in vitro* DNA studies. A glycated DNA can be identified by comparing it with the native DNA and determining the modification, mutation, or nicking of DNA (Ahmad et al. 2017).

Hypothesizing that oxidative stress is one of the major contributors to glycation and protein aggregation, the superfood *Spirulina*'s species, *Arthrospira platensis*, was selected for the test owing to its high antioxidant potential. In this study, *A. platensis* was sonicated, and its aqueous extract was used to determine its phytochemical constituents, antioxidant properties, and ability to prevent glycation and protein aggregation. Our results indicate that the extract could effectively reduce the structural and functional changes in HB as induced by three sugars, namely glucose, fructose, and ribose. Furthermore, using pBR322 as a model for DNA, an *A. platensis* aqueous extract was proven to be efficient in preventing its glycooxidation.

Materials and Methods

Materials

Axenic culture of *A. platensis* PCC 7345 was procured from Dr. Vani, BITS Pilani (Hyderabad, India). BG11 medium was purchased from Himedia Laboratories Pvt. Ltd., Mumbai, India. All other chemicals were purchased from SD Fine-Chem. Ltd., Mumbai, India.

Preparation of Extract

The exponential phase, that is, 11–15-day-old culture of *A. platensis* PCC 7345, was collected and centrifuged at 10,000 rpm for 10 mins at 4°C. The biomass was collected and washed twice with distilled water (D/W). The biomass was dissolved in D/W at 100 mg/mL concentration and sonicated at 37°C for 10 min at 15 pulses/min. This was further centrifuged, and the supernatant was used for further analysis.

Phytochemical Screening of the Extract

Quantification of Biochemical Compounds

The carbohydrate and protein content of the samples were quantified by 3,5-dinitrosalicylic acid (DNSA) and Bradford's method at 540 nm and 595 nm, respectively, using a spectrophotometer. Total phenolic content was determined using a modified Folin-Ciocalteu's method compared with gallic acid as the standard (Singh et al., 2002). Total flavonoid content was estimated using the aluminum chloride method, representing the result as quercetin equivalents/ml (Kumar et al., 2022).

Antioxidant Activity Profiling

The free radical scavenging activity of the extract was determined by the 2, 2-diphenyl-1-picrylhydrazyl (DPPH) assay. The Griess reagent was freshly prepared and used for the nitric oxide radical scavenging assay. Five milliliters of a varying concentration of extract (20–100 μ g/mL) was mixed with an overnight prepared 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) solution, and readings were taken after 30 min of incubation at 734 nm (Rajurkar & Hande, 2011). The percent inhibition in all three assays was calculated using the following formula:

$$\% \text{ Inhibition} = [(\text{Abs Control} - \text{Abs Extracts}) / \text{Abs Control}] \times 100.$$

Determination of the Presence of Phycobiliproteins in the Extract

The cyanobacterium *A. platensis* contains three phycobiliproteins, namely, C-phycocyanin (C-PC), allophycocyanin (APC), and phycoerythrin (PE). All the light-harvesting bilin proteins were determined fluorometrically using a Cary Eclipse Fluorescence Spectrophotometer. The excitation wavelengths of C-PC, APC, and PE are 620, 651, and 566 nm, respectively,

and the emission was recorded at 500–750 nm (Dagnino-Leone et al., 2022).

Incubation for Antiglycating Assays

A. platensis aqueous extract (100 µg/mL) was tested for its antiglycation potential using HB (1 mg/mL) as the protein and various sugars, namely, glucose (100 mg/mL), fructose (100 mg/mL), and ribose (10 mg/mL). Glycated control contained sugar and protein, sodium phosphate buffer (0.1 M, pH 7.4), and sodium azide (30 mM), while the test sample contained glycated control with the extract. The tubes were incubated at 37°C for 28 days and then subjected to antiglycation assays immediately (Banan & Ali, 2016).

Determination of Early and Late Glycated Products

Browning

The extent of browning in the incubated samples was measured at 420 nm using the Shimadzu UV 1800 spectrophotometer. The results of test samples were compared with relative percentages concerning the standard (Banan & Ali, 2016).

Fructosamine and HbA_{1c}

The ketoamine or Amadori product, formed during the initial stages of glycation, was determined using a nitroblue tetrazolium (NBT) assay. One hundred microliters of NBT reagent was incubated with 10 µL of glycated samples at 37°C for 15 min. 0.89 mL of D/W was added to this reaction mixture to increase the volume to 1 mL. Absorbance was measured at 530 nm, and the values were calculated in µmol/L using 1-deoxy-1-morpholinofructose (DMF) as a standard.

The amount of fructosamine and HbA_{1c} are highly correlated. According to the ADA, 292 mmol/l of serum fructosamine content is equivalent to 6.57% of HbA_{1c} (Cohen et al., 2003). Thus, using the fructosamine content, HbA_{1c} levels in incubated samples were determined using the following formula (Vasista et al., 2015):

$$\text{HbA}_{1c} = 0.017 \times \text{fructosamine level} (\mu\text{mol/L}) + 1.61.$$

Protein Carbonyl Content

The carbonyl content in the glycated samples was measured using the 2,4-dinitrophenylhydrazine (DNPH) assay. One hundred microliters of the incubated samples were added to 400 µL of DNPH reagent and incubated in the dark for an hour. 500 µL of trichloroacetic acid (TCA) was added to precipitate the protein in the samples. After 5 min of incubation on ice, the samples were centrifuged at 4°C for 10

min at 10,000 rpm. The pellet was collected and washed three times with ethanol:ethyl acetate (1:1) before suspending in 6M guanidine hydrochloride. The solutions were diluted up to 1 mL using D/W, and the absorbance was determined spectrophotometrically at 370 nm using a Shimadzu UV 1800 spectrophotometer (Banan & Ali, 2016).

Total Fluorescent AGEs

The incubated samples were diluted ten times using D/W and checked for fluorescence by excitation wavelength at 370 nm and emission at 440 nm (Banan & Ali, 2016; Thornalley & Rabbani, 2014).

Fluorescent Individual AGEs

Some of the AGEs have been known to exhibit fluorescence. The incubated samples were analyzed fluorometrically for the same and checked for the formation of fluorescent AGEs (Thornalley & Rabbani, 2014). The fluorescent AGEs mentioned in Table 1 were determined.

Determination of Band Shift of Heme Proteins in HB

All heme proteins exhibit a Soret peak in the blue wavelength region of the visible spectrum. The red band shift in the glycated proteins and its prevention were determined spectrophotometrically between 380 and 450 nm (Behroozi et al., 2014).

Analysis of Glycation-induced Protein Aggregation

Protein Aggregation Index

The presence of protein aggregates in the sample was determined spectrophotometrically, and the protein aggregation index was then further calculated using the following formula:

$$\text{Aggregation index} = [A_{340} / (A_{280} - A_{340})] \times 100.$$

Table 1. Individual Fluorescent AGEs with Excitation and Emission Wavelengths.

Sr. No.	Fluorescent AGEs	Excitation Wavelength (nm)	Emission Wavelength (nm)
1.	Total AGEs	350	440
2.	Argpyrimidine	320	380
3.	Vesperlysine A (LMI)	350	405
4.	Pentosidine	335	385
5.	Crossline	380	440

Source: Adapted from Thornalley and Rabbani (2014).

Abbreviation: AGE, advanced glycation end product.

Protein Aggregation by Thioflavin-T

AGEs trigger the formation of protein aggregates. These increased β -structures were determined using a fluorescent dye, Thioflavin-T (ThT). The incubated sample was diluted 10 times with D/W. 40 μ L of 8 mM ThT in 100 mM of sodium phosphate buffer (pH 7.4) was added to 1.0 mL of diluted sample and kept for incubation for 15 min at RT. The readings were acquired using 444/485 nm excitation/emission filters on a Cary Eclipse Fluorescence Spectrophotometer (Banan & Ali, 2016).

Prevention of Glyoxidative Damage to DNA

In vitro glycation of plasmid DNA in the presence of *A. platensis* extract:

pBR322 plasmid DNA (0.25 μ g) in 100 mM potassium phosphate buffer (pH 7.4) was incubated with lysine (20 mM), MG (20 mM), and FeCl₃ (100 μ M) in the presence and absence of *A. platensis* extract. The reaction mixtures of samples were incubated at 37°C for 2 h. pBR322 plasmid DNA (0.25 μ g) in 100 mM potassium phosphate buffer (pH 7.4) without a glycation system, and *A. platensis* extract was used as a control (Kumar et al., 2022).

Agarose Gel Electrophoresis of a Glycated Plasmid DNA Sample

Ten microliters of incubated samples were mixed with 2 μ L of 6X gel loading dye and loaded onto a 1% agarose gel prepared in TAE buffer. Electrophoresis was carried out at 85 V, and the gel was run up to two-third of its length. The gel was stained with fluorescent dye Ethidium Bromide (5 μ g/ml) and visualized using Gel-Doc.

Determination of Change in Band IntDen

DNA images from Gel-Doc were analyzed using ImageJ (Version 1.53s) to measure the integrated density (IntDen). The colors of the gel image were inverted/converted into dark bands-light background. A graph of gray area vs. density was plotted using the software. The IntDen was further calculated, and an increase/decrease in bands was determined.

Statistical Analysis

All the graphs were plotted and analyzed using GraphPad Prism version 9.4.0. Statistical significance of the data was evaluated using ordinary one-way ANOVA, and the Brown-Forsythe test was applied wherever applicable.

Table 2. Preliminary Biochemical Screening of *A. platensis* Aqueous Extract.

Concentration of Extract (μ g/mL)	Carbohydrate Content (mg/mL)	Protein Content (mg/mL)	Total Phenolic Content (Gallic Acid Equivalents/mL)	Total Flavonoid Content (Quercetin Equivalents/mL)
20	0.04	6.2137	0.0980	0.1726
40	0.132	7.5496	0.1349	0.257
60	0.162	8.3129	0.2199	0.7482
80	0.24	13.0458	0.3184	1.0239
100	0.32	14.1526	0.3851	1.8393




Figure 1. Antioxidant Activities of *A. platensis* Aqueous Extract.

Note: Results expressed in the percentage of inhibition; mean ($n = 5$).

Abbreviations: DPPH, 2,2-diphenyl-1-picrylhydrazyl; ABTS, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid).

p -values < 0.05 were considered statistically significant. The significance was further indicated in the graph as follows: ** $p < 0.05$; *** $p < 0.005$; **** $p < 0.0005$.

Results

Biochemical Screening of the Extract

Aqueous extracts prepared from biomass at concentrations of 20–100 $\mu\text{g/mL}$ were subjected to biochemical screening. There was a linear increase in all the measured phytoconstituents. A comparison of carbohydrate and protein content in the biomass clearly indicates the high level of protein present in the cells (Table 2). Similarly, the flavonoid content was found to be almost five times as high as the phenolic acid content. These compounds have been known to impart antioxidant properties to the organism.

Antioxidant Profiling of the Extract

DPPH assay, or free radical scavenging activity assay, and ABTS decolorization assay help in the determination of the total antioxidant potential of the sample, herein the extract (Mahdi-Pour et al., 2012). However, the assays are performed in different solvents, that is, methanol and D/W, respectively (Figure 1).

Thus, the higher solubility of organic compounds in organic solvents could be one of the reasons for the difference in the IC_{50} values. The IC_{50} values were calculated from the equations obtained through graphs. IC_{50} values of DPPH, NO radical, and ABTS scavenging assays were 42.5, 52.29, and 31.09 $\mu\text{g/mL}$, respectively (Table 3).

Determination of the Presence of Phycobiliproteins

A. platensis contains a phycobilisome that captures light energy and helps the organism do photosynthesis. Three phycobiliproteins are present in this complex, which was

determined by fluorometer. The maximum PE, APC, and C-PC emission was observed at 565, 651, and 620 nm (Figure 2).

Quantification of Glycated Products

Browning

Browning is an indicator of the Maillard reaction, which is the first stage of glycation. Considering HB + sugars as the standard (100%), the percent reduction of browning of the samples was determined. The highest reduction of 53.83% was observed in glucose, followed by fructose and ribose, where the extract could inhibit browning by 42.62% and 37.43%, respectively (Figure 3).

Protein Carbonyl Content

Protein carbonyls are formed during the second glycation stage, which further forms AGEs. The aqueous extract was found most effective by reducing 65.14% of the protein carbonyl content in the presence of ribose, followed by fructose, where the reduction was 36.17% (Figure 4). Though a similar reduction (21.74%) was also observed in HB incubated with glucose and the sample, the result was insignificant.

Fructosamine Content and HbA_{1c} Level

A standard graph of DMF was plotted, and values of fructosamine were extrapolated from the graph. The HbA_{1c} level was further calculated using the previously mentioned formula (Table 4). In the presence of the extract, both fructosamine and HbA_{1c} levels were reduced in all three sugars.

Total Fluorescent AGE

Some AGEs exhibit fluorescent properties. Inhibition was observed in all glycated samples (Figure 5) with extracts compared with the standards (HB + sugar).

Table 3. Antioxidant Assays.

Concentration ($\mu\text{g/mL}$)	Free Radical Scavenging Assay (Reduction Percentage)		Nitric Oxide Radical Scavenging Assay (Reduction Percentage)		ABTS Radical Scavenging Assay (Reduction Percentage)	
	Standard	Extract	Standard	Extract	Standard	Extract
20	27.91	49.17	39.54	57.14	26.97	20.35
40	36.51	52.80	52.01	69.84	33.07	27.77
60	42.68	60.23	59.88	73.75	45.37	38.45
80	49.24	68.31	72.26	80	55.67	47.49
100	60.96	80.52	75.64	82.43	76.55	60.55
Equation from graph	$y = 0.3942x + 19.811$	$y = 0.3911x + 38.743$	$y = 0.4622x + 32.138$	$y = 0.3037x + 54.41$	$y = 0.6088x + 10.998$	$y = 0.5006x + 8.886$
R^2 value	0.9858	0.9633	0.9743	0.9239	0.9596	0.9918
IC_{50} value ($\mu\text{g/mL}$)	76.58	45.97	38.65	52.29	64.063	31.09

Abbreviation: ABTS, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid).




Figure 2. Fluorescence Spectrum of Phycobiliproteins.

Note: Results expressed arbitrary units (a.u.) of mean values ($n = 5$).

Abbreviations: PE, phycoerythrin; APC, allophycocyanin; CPC, C-phycoerythrin.




Figure 4. Measurement of Carbonyl Content.

Note: Results expressed reduction percentage $\pm$ standard deviation ($n = 8$); $p < 0.005$.

Abbreviations: HB, hemoglobin; G, glucose; F, fructose; R, ribose; S, sample, i.e., *A. platensis* aqueous extract.




Figure 3. Reduction of Browning Using Different Hemoglobin and Sugar Mixtures.

Note: Results expressed reduction percentage $\pm$ standard deviation ($n = 8$); $p < 0.005$.

Abbreviations: HB, hemoglobin; G, glucose; F, fructose; R, ribose; S, sample, i.e., *A. platensis* aqueous extract.




Figure 5. Determination of Total Fluorescent AGEs in Glycated Sample.

Note: Results expressed in arbitrary units (a.u.) of mean values ($n = 8$); $p < 0.05$.

Abbreviations: AGE, advanced glycation end product; HB, hemoglobin; G, glucose; F, fructose; R, ribose; S, sample, i.e., *A. platensis* aqueous extract.

Table 4. Fructosamine and HbA_{1c} Content in Glycated Sample.

	Fructosamine ($\mu\text{mol/L}$)	HbA _{1c}
Hemoglobin	0.1674	1.638458
Hemoglobin + Extract	0.2604	1.654268
Hemoglobin Glucose	1.5438	1.872446
Hemoglobin + Glucose+ Extract	0.3906	1.676402
Hemoglobin + Fructose	1.4322	1.853474
Hemoglobin + Fructose + Extract	0.3906	1.676402
Hemoglobin + Ribose	1.767	1.91039
Hemoglobin + Ribose + Extract	0.744	1.73648

Note: Results expressed in $\mu\text{mol/L}$ of fructosamine and HbA_{1c} level $\pm$ standard deviation ($n = 8$); $p < 0.005$.

Abbreviation: HbA_{1c}, glycated hemoglobin.

Determination of Individual AGEs

Kiviat diagrams/radar charts were used to represent the different AGEs present in the incubated sample and their respective reductions in the presence of *A. platensis* aqueous extract. The blue line indicates the level of fluorescent AGEs in arbitrary units as a positive standard, that is, HB + sugar, while the red line indicates the level of AGEs after treatment with extract. All the sugars could induce all five types of fluorescent AGEs in HbA_{1c} (Figure 6).

It can be observed from the graphical representation that glucose could induce the highest number of total AGEs, followed by crossline and vesper lysine A. Although *A. platensis* extract could reduce the fluorescent intensity of all AGEs, it was found more effective in reducing total AGEs and crosslines. Similarly, the extract could effectively reduce

total AGEs in fructose-mediated glycation of HB, while in the case of ribose-mediated glycation, extract was more effective in reducing pentosidine and argpyrimidine.

Soret-Band Statue of Heme Proteins

A decrease in Soret peak as well as redshift can be observed in HbA_{1c} (Figure 7). The highest change has been observed in HB incubated with ribose, fructose, and glucose, respectively. The extract had almost no effect on the absorptivity of heme proteins and could effectively preserve the absorptivity in the presence of all three sugars.

Quantification of Glycation-induced Aggregation of Proteins

Protein Aggregation Index

The highest aggregation index was observed in HB + glucose, followed by fructose and ribose. The aqueous extract of *A. platensis* could effectively decrease the aggregation of protein in all three sugars. The most significant decrease was observed in the case of glucose (Figure 8).

Protein Aggregation by ThT

A structural alteration in glycated proteins leads to an increase in amyloid fibrils and cross β -structure. These changes were monitored using the fluorescent dye ThT (Figure 9). A significant decrease in β -structure can be observed when HB + sugars were incubated with extracts in comparison with their standard.

Effect of *A. platensis* Extract on DNA Damage

Plasmid DNA pBR322 alone (lane 1) displayed two bands, band 1 indicating a linear form, and the lower band, that is, band 2, indicating supercoiled form (Figure 10a).

Glycation causes strand breaks in the supercoiled form, thereby turning them linear as observed in lane 2 (pBR322 + glycation system). Lane 3 containing DNA and *A. platensis* extract, appeared similar to lane 1 implying that the extract did not affect the DNA. Lane 4 containing DNA + glycation system + extract seemed to be similar to lane 1, with lesser intensity in the linear band (Figure 10b) and higher intensity in the supercoiled form (Figure 10c). This indicates that the




Figure 6. Kiviatt Diagram Representing the Formation of Individual AGEs in the Presence of HB + Sugars. (a) Glucose, (b) Fructose, and (c) Ribose.

Note: All the axes representing each AGEs originate from the same center, and the difference between the blue and red line is directly proportional to the change in the level of AGEs.

Abbreviations: AGE, advanced glycation end product; HB, hemoglobin; G, glucose; F, fructose; R, ribose; S, sample, i.e., *A. platensis* aqueous extract.




Figure 7. UV-vis Spectrum of Glycated Proteins.

Note: Results expressed in arbitrary units (a.u.) of mean values ($n = 8$); $p < 0.05$.

Abbreviations: HB, hemoglobin; G, glucose; F, fructose; R, ribose; S, sample, i.e., *A. platensis* aqueous extract.




Figure 8. Determination of Aggregation Index of Protein.

Note: Results expressed in arbitrary units (a.u.) $\pm$ standard deviation ($n = 8$); $p < 0.05$.

Abbreviations: HB, hemoglobin; G, glucose; F, fructose; R, ribose.




Figure 9. Determination of Protein Aggregation by Thioflavin T-Graph of Intensity vs. Wavelength.

Note: Results expressed in arbitrary units (a.u.) of mean values ($n = 3$); $p < 0.05$.

Abbreviations: HB, hemoglobin; G, glucose; F, fructose; R, ribose; S, sample, i.e., *A. platensis* aqueous extract; ThT, Thioflavin-T dye.




Figure 10. Glycooxidative DNA Damage and Its Subsequent Prevention by *A. platensis* Aqueous Extract.

(a) Lanes: 1 – DNA alone; 2 – DNA + Glycation System (MG + FeCl₃ + Lys); 3 – DNA + *A. platensis* Aqueous Extract; 4 – DNA + Glycation System + Extract [Incubation Time: 180 min; Stained with Ethidium Bromide].

(b) Graph of Gray Value vs. Distance ~ for Band 1, i.e., Linear Form of DNA as Analyzed by ImageJ Lanes as Described in (a).

(c) Graph of Gray Value vs. Distance ~ for Band 2, i.e., Supercoiled Form of DNA as Analyzed by ImageJ Lanes as Described in (a).

Abbreviations: MG, methyl glyoxal; Lys, lysine.

Table 5. Comparison of IntDen between DNA Lanes in Band 1 (Linear Form) and Band 2 (Supercoiled Form).

Band 1			Band 2		
Lane	IntDen	Percentage Increase/Decrease	Lane	IntDen	Percentage Increase/Decrease
1	5437.711	0	1	17473.309	0
2	24498.007	+350.52	2	401.678	−97.70
3	4235.437	−22.11	3	23004.986	+31.66
4	672.799	−87.63	4	19986.744	+14.38

Note: Lanes: 1 – DNA alone; 2 – DNA + Glycation System (MG + FeCl₃ + Lys); 3 – DNA + *A. platensis* Aqueous Extract; 4 – DNA + Glycation System + Extract. The increase and decrease have been indicated with + and − signs before the value, respectively.

extract could reverse the strand break induced by the glycation system on the DNA or prevent the same. These results were further analyzed using ImageJ, and it can be observed that MG could convert almost 97% of DNA from supercoiled form to linear form (Table 5). However, the presence of the extract was found to inhibit the perturbation in DNA and repair it back to its native supercoiled form.

Discussion

Glycation is a spontaneous, non-enzymatic process occurring in hyperglycemia. Excess sugar in the body starts reacting

with the biomolecules and altering their structural integrity, compromising their functionality. The current research is focused on detecting early glycation products and AGEs formed by HB in reaction with the three most abundant sugars in the body, that is, glucose, fructose, and ribose. Since ribose was the most highly reactive sugar among the three, glycation products appeared to be higher in comparison with fructose and glucose.

The first step of glycation is the Maillard reaction, wherein the carbonyl group from reducing sugars reacts with free amino groups to form a reversible Schiff base. It further undergoes autooxidation to form ketamine (e.g.,

fructosamine). Reactive protein carbonyl intermediates trigger the formation of AGEs and protein aggregates. All three stages of glycation liberate hydroxyl radicals, reactive oxygen species (ROS), reactive nitrogen species (RNS), and superoxide radicals in the presence of transition metals, which in turn leads to acceleration of the glycation process (Smith & Thornalley, 1992; Suantawee et al., 2015).

As oxidative stress is one of the triggering factors in the manifestation of glycation (Paramanya & Ali, 2019), *A. platensis*, a source of a powerful antioxidant, was checked for its antiglycating property. Fluorometric analysis of the extract shows the presence of all three phycobiliproteins, namely, APC, C-PC, and PE, all of which have been known to contain antioxidant properties. Notably, the extract was found to be more efficient in preventing and/or reversing the damage done to HB than glucose followed by fructose and ribose consecutively. It can be seen from our results that the extract is more efficient at preventing the formation of AGEs than it is at inhibiting the formation of early glycation products. One of the reasons for this could be narrowed down to the faster reaction rates in the formation of early glycation products with respect to AGE. The determination of bioactive compounds in the extract and improving its site-specificity might help prevent the glycation of hemoglobin at earlier stages.

The data collected in this research also represent the formation of different types of fluorescent AGEs induced by all three sugars in HB. The aqueous extract of *A. platensis* could reduce the formation of these AGEs; however, there was no direct similarity in reduction of AGEs when compared with HbA_{1c} formed by the three sugars. Behroozi and colleagues (2014) reported that glycation of HB caused a red shift of the heme proteins. The findings of this research paper coincide with this study, wherein the changes in decreased absorptivity of heme proteins were attenuated effectively by *A. platensis* extract.

Ribose, the most reactive of the three tested sugars, appeared to elevate the levels of early and late glycation products and protein aggregates. Thus, the glycation potential of the sugars can be stated as ribose>fructose>glucose. The antiglycating potential of the extract was thus highest in glucose-mediated glycation, followed by fructose and ribose. It is noteworthy that the extracts were found to be more effective in preventing glycations at early stages compared to the reduction observed in the late stages and during protein aggregation.

The final part of the study dealt with determining the perturbation of DNA using the pBR322 model and further prevention of strand breaks using extract. MG, a dicarbonyl sugar, was used to induce strand breaks and convert DNA from supercoiled to linear (Ahmad et al., 2017; Jha et al., 2018). However, the *A. platensis* extract was found coherent, decreasing linearity and restoring the DNA back to its supercoiled form. Although *A. platensis* has been previously

shown to have an anti-DNA damage effect in cadmium chloride-induced DNA damage (Aly et al., 2018), this article is the first to report DNA protective effects in glycoxidative DNA damage.

It has to be noted that this study has several limitations, which include the use of high sugar concentrations to elevate the glycation process. It is still unclear exactly how the compounds in the extract get metabolized *in vivo* and the plethora of factors orchestrating this process, like the amount and extent of glycemia. Based on the results, it can be concluded that the antioxidant properties of *A. platensis* could be one of the major contributing factors. However, a comprehensive study of the extract could help determine the bioactive compound involved in contributing to the antiglycating property of *A. platensis*.

Conclusion

A. platensis aqueous extract contains all three phycobiliprotein, which possess antioxidant properties. The current study on HB shows that this extract could effectively attenuate the damages occurring due to glycation induced by the three tested sugars, namely, glucose, fructose, and ribose. The extract could also prevent glycoxidative damage on pBR322, implying that *A. platensis* can be considered to possess antiglycating properties.

Summary

Spirulina (*A. platensis*) extract was evaluated for its antiglycating potential against three sugars, namely, glucose, fructose, and ribose. For protein, HB was used, and it was found that the extract prevented both early and late stages of glycation and protein aggregation against all three sugars. The extract also prevented the DNA strand breaks in pBR322 caused by glycoxidation.

Acknowledgments

We gratefully acknowledge Dr. Vani from BITS Pilani, Hyderabad, for providing an axenic culture of *A. platensis* PCC 7345. The authors would also like to extend their sincere appreciation to the Researchers Supporting Project number (RSP2023R154), King Saud University, Riyadh, Saudi Arabia.

Abbreviations

ABTS: 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid); ADA: American Diabetes Association; AGEs: advanced glycation end products; BSA: bovine serum albumin; DMF: 1-deoxy-1-morpholinofructose; DNPH: 2,4-dinitrophenylhydrazine; DNSA: 3,5-dinitrosalicylic acid; DPPH: 2,2-diphenyl-1-picrylhydrazyl; F: fructose; G: glucose; HB: hemoglobin; HbA_{1c}: glycated hemoglobin; Lys: lysine; MG: methylglyoxal; NBT: nitroblue tetrazolium; R: ribose; RNS: reactive nitrogen species; ROS: reactive oxygen species; S: sample, i.e., *A. platensis* PCC 7345 aqueous extract (100 µg/ml); TCA: trichloroacetic acid; ThT: thioflavin-T dye.

Authors' Contributions

Additiya Paramanya: Methodology, Formal analysis, Investigation, and Writing—original draft. Mohammad Abul Farah and Khalid Mashay Al-Anazi: Conceptualization, Funding acquisition, Writing—review & editing, and Formal analysis. Hari Prasad Devkota: Writing—review & editing and Formal analysis. Ahmad Ali: Conceptualization, Writing—review & editing, Supervision, Project administration, and Validation.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical Approval

Not Applicable.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was funded by the Researchers Supporting Project number (RSP2023R154), King Saud University, Riyadh, Saudi Arabia.

Informed Consent

Not Applicable.

ORCID iD

Ahmad Ali  <https://orcid.org/0000-0003-4467-5387>

References

- Ahmad, R., Sah, A. K., & Ahsan, H. (2017). Biochemistry and pathophysiology of glycation of DNA: Implications in diabetes. *Archives of Clinical and Biomedical Research*, 1(1), 32–47.
- Ahmad, S., Moinuddin, Shahab, U., Habib, S., Khan, M. S., Alam, K., & Ali, A. (2014). Glycoxidative damage to human DNA: Neo-antigenic epitopes on DNA molecule could be a possible reason for autoimmune response in type 1 diabetes. *Glycobiology*, 24(3), 281–291. <https://doi.org/10.1093/glycob/cwt109>
- Aly, F. M., Kotb, A. M., & Hammad, S. (2018). Effects of *Spirulina platensis* on DNA damage and chromosomal aberration against cadmium chloride-induced genotoxicity in rats. *Environmental Science and Pollution Research International*, 25(11), 10829–10836. <https://doi.org/10.1007/s11356-018-1329-3>
- American Diabetes Association (2022). Standards of medical care in Diabetes-2022 abridged for primary care providers. *Clinical Diabetes*, 40(1), 10–38. <https://doi.org/10.2337/cd22-as01>
- Banan, P., & Ali, A. (2016). Preventive effect of phenolic acids on *in vitro* glycation. *Annals of Phytomedicine*, 5, 97–102.
- Behroozi, J., Divsalar, A., & Saboury, A. A. (2014). Honey bee venom decreases the complications of diabetes by preventing hemoglobin glycation. *Journal of Molecular Liquids*, 199, 371–375. DOI: 10.1016/j.molliq.2014.09.034.
- Bunn, H. F., Haney, D. N., Gabbay, K. H., & Gallop, P. M. (1975). Further identification of the nature and linkage of the carbohydrate in hemoglobin A1c. *Biochemical and Biophysical Research Communications*, 67(1), 103–109. [https://doi.org/10.1016/0006-291x\(75\)90289-2](https://doi.org/10.1016/0006-291x(75)90289-2)
- Cohen, R. M., Holmes, Y. R., Chenier, T. C., & Joiner, C. H. (2003). Discordance between HbA_{1c} and fructosamine: Evidence for a glycosylation gap and its relation to diabetic nephropathy. *Diabetes Care*, 26(1), 163–167. <https://doi.org/10.2337/diabetes.26.1.163>
- Dagnino-Leone, J., Figueroa, C. P., Castañeda, M. L., Youton, A. D., Vallejos-Almirall, A., Agurto-Muñoz, A., Pavón Pérez, J., & Agurto-Muñoz, C. (2022). Phycobiliproteins: Structural aspects, functional characteristics, and biotechnological perspectives. *Computational and Structural Biotechnology Journal*, 20, 1506–1527. <https://doi.org/10.1016/j.csbj.2022.02.016>
- Jha, P., Momin, A. R., Kumar, D., & Ali, A. (2018). Reversal of glycoxidative damage of DNA and protein by antioxidants. *Annals of Phytomedicine*, 7(1), 1–5.
- Kumar, D., Shah, N., Al-Anazi, K. M., Farah, M. A., Ali, M. A., Lee, J., & Ali, A. (2022). Potential role of *Tribulus terrestris* fruit extract in inhibition of advanced glycation end products. *Pharmacognosy Magazine*, 18(78), 341.
- Mahdi-Pour, B., Jothy, S. L., Latha, L. Y., Chen, Y., & Sasidharan, S. (2012). Antioxidant activity of methanol extracts of different parts of *Lantana camara*. *Asian Pacific Journal of Tropical Biomedicine*, 2(12), 960–965. [https://doi.org/10.1016/S2221-1691\(13\)60007-6](https://doi.org/10.1016/S2221-1691(13)60007-6)
- Paramanya, A., & Ali, A. (2019). Role of oxidative stress in Biological systems. *Middle East Journal of Science*, 5 (2), 155–162. DOI:10.23884/mejs.2019.5.2.07.
- Rajurkar, N. S., & Hande, S. M. (2011). Estimation of phytochemical content and antioxidant activity of some selected traditional Indian medicinal plants. *Indian Journal of Pharmaceutical Sciences*, 73(2), 146–151. <https://doi.org/10.4103/0250-474x.91574>
- Singh, R. P., Murthy, K. N. C., & Jayaprakasha, G. K. (2002). Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using *in vitro* models. *Journal of Agricultural and Food Chemistry*, 50(1), 81–86. <https://doi.org/10.1021/jf010865b>
- Singh, V. P., Bali, A., Singh, N., & Jaggi, A. S. (2014). Advanced glycation end products and diabetic complications. *The Korean Journal of Physiology & Pharmacology*, 18(1), 1.
- Smith, P. R., & Thornalley, P. J. (1992). Mechanism of the degradation of non-enzymatically glycated proteins under physiological conditions. Studies with the model fructosamine, N epsilon-(1-deoxy-D-fructos-1-yl)hippuryl-lysine. *European Journal of Biochemistry*, 210(3), 729–739. <https://doi.org/10.1111/j.1432-1033.1992.tb17474.x>
- Suantawee, T., Wesarachanon, K., Anantsuphasak, K., Daenphetploy, T., Thien-Ngern, S., Thilavech, T., Pasukamonset, P., Ngamukote, S., & Adisakwattana, S. (2015). Protein glycation inhibitory activity and antioxidant capacity of clove extract. *Journal of Food Science and Technology*, 52(6), 3843–3850. <https://doi.org/10.1007/s13197-014-1452-1>
- Thornalley, P. J., & Rabbani, N. (2014). Detection of oxidized and glycated proteins in clinical samples using mass spectrometry—a user's perspective. *Biochimica et Biophysica Acta*, 1840(2), 818–829. <https://doi.org/10.1016/j.bbagen.2013.03.025>
- Vasista, P., Tziaferi, V., Greening, J., & Sundaram, P. (2015). HbA_{1c}: Is it a reliable measure of glycaemic control in all patients with type 1 diabetes mellitus? In *Endocrine abstracts*. Vol. 39. Bioscientifica. DOI: 10.1530/endoabs.39.EP49.